



FUTIFMS International Conference

Weighted Hardy Inequalities for Singular Elliptic Problems: Theory, Applications, and Computational Perspectives

O. P. Durojaye

Department of Mathematics, Federal University of Technology Ilaro (FUTI),
Ilaro, Ogun State, Nigeria.

Corresponding author:
Email: opidi2000@gmail.com

Abstract Weighted Hardy inequalities constitute a cornerstone of modern functional analysis and the theory of partial differential equations (PDEs). By extending the classical Hardy inequality through the incorporation of weight functions, these inequalities provide a robust framework for handling singularities inherent in elliptic boundary value problems. Their applications are vast, encompassing Sobolev space theory, variational methods, mathematical physics, and numerical approximation. This paper reviews the theoretical foundations of weighted Hardy inequalities, emphasizing their critical role in establishing coercivity, stability, and a priori estimates for singular elliptic problems. Furthermore, we examine their computational implications for finite element methods (FEM) and adaptive numerical schemes. Finally, we highlight future research directions, including fractional Hardy inequalities and data-driven numerical methods.

Keywords: Hardy inequality; weighted Hardy inequality; Sobolev spaces; elliptic PDEs; finite element method; numerical analysis.

1. Introduction

Functional inequalities, which bound the norms of functions by the norms of their derivatives, are indispensable tools in the analysis of differential equations. They are essential for proving the existence, uniqueness, regularity, and stability of solutions [4, 6]. Among these, the Hardy inequality stands out as a fundamental result. Originally formulated in the early 20th century, it has since been generalized to weighted, fractional, and Riemannian manifold settings [1, 2].

In the context of partial differential equations, standard Sobolev embeddings often fail to control functions with strong singularities. Weighted Hardy inequalities bridge this gap by incorporating specific weight functions—often related to the distance to the boundary or the distance to a singular point—allowing for the rigorous treatment of singular potentials and boundary singularities [3, 7]. This paper synthesizes the theoretical underpinnings of these inequalities and

explores their profound impact on both the continuous analysis and the discrete numerical approximation of singular elliptic problems.

2. Mathematical Preliminaries

Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with a sufficiently regular boundary $\partial\Omega$. The foundation of our analysis rests on Sobolev spaces. The space $W^{1,p}(\Omega)$ for $1 \leq p \leq \infty$ consists of functions in $L^p(\Omega)$ whose weak first-order derivatives also belong to $L^p(\Omega)$ [4, 8]. The subspace $W_0^{1,p}(\Omega)$ is defined as the closure of $C_0^\infty(\Omega)$ (smooth functions with compact support) in the $W^{1,p}$ norm.

Classical Hardy Inequality:

In one dimension, for any $u \in C_0^\infty(0, \infty)$, the classical Hardy inequality states that:

$$\int_0^\infty \frac{|u(x)|^2}{x^2} dx \leq 4 \int_0^\infty |u'(x)|^2 dx \quad (1)$$

The constant 4 is optimal and cannot be improved [1, 2]. In n dimensions ($n \geq 3$), for $u \in C_0^\infty(\mathbb{R}^n)$, the inequality takes the form:

$$\int_{\mathbb{R}^n} \frac{|u(x)|^2}{|x|^2} dx \leq \frac{4(n-2)}{2} \int_{\mathbb{R}^n} |\nabla u(x)|^2 dx \quad (2)$$

Weighted Hardy Inequality:

The weighted generalization introduces positive weight functions $w(x)$ and $v(x)$. The inequality asserts that:

$$\int_\Omega w(x) |u(x)|^p dx \leq C \int_\Omega v(x) |\nabla u(x)|^p dx \text{ for all } u \in W_0^{1,p}(\Omega), \quad (3)$$

where $C > 0$ is a constant independent of u . The validity of this inequality depends heavily on the Muckenhoupt A_p conditions or specific structural assumptions on the weights w and v [5].

3. Main Results

The most prominent application of Hardy-type inequalities in PDEs involves singular potentials. A standard weighted Hardy inequality for bounded domains Ω containing the origin, with $1 < p < n$, states that there exists a constant $C = C(n, p, \Omega) > 0$ such that:

$$\int_\Omega \frac{|u(x)|^p}{|x|^p} dx \leq C \int_\Omega |\nabla u(x)|^p dx \quad (4)$$

for all $u \in W_0^{1,p}(\Omega)$ [3, 4]. The proof typically relies on integration by parts, Hölder's inequality, and careful manipulation of radial vector fields.

When the singularity is located on the boundary rather than in the interior, the weight function is replaced by the distance to the boundary, $d(x) = \text{dist}(x, \partial\Omega)$. For domains with smooth boundaries, the Hardy inequality takes the form:

$$\int_{\Omega} \frac{|u(x)|^p}{d(x)^p} dx \leq C \int_{\Omega} |\nabla u(x)|^p dx \quad (5)$$

This boundary-weighted version is crucial for analyzing problems with boundary degeneracies. Furthermore, in the critical case where $p=n$, the standard Hardy inequality fails, and logarithmic corrections or Sobolev-type embeddings must be employed to recover coercivity [3, 5].

4. Applications

The utility of weighted Hardy inequalities extends across several domains of mathematical analysis and scientific computing.

4.1. Coercivity and Singular Elliptic Problems

Consider the singular elliptic operator $L = -\Delta - \frac{\mu}{|x|^2}$, where $\mu > 0$ is a parameter. The associated energy functional involves the term

$\int |\nabla u|^2 - \mu \int \frac{u^2}{|x|^2}$. The Hardy inequality guarantees that if $\mu < \frac{(n-2)^2}{4}$ (the optimal Hardy constant), the operator L remains strictly coercive on $H_0^1(\Omega)$ [7]. This coercivity is the linchpin for applying the Lax-Milgram theorem to establish the existence and uniqueness of weak solutions.

4.2. Variational Methods and Stability

In variational formulations, Hardy inequalities provide essential a priori estimates. They allow analysts to control the L^p norm of a function by a singular weight using only the L^p norm of its gradient [6]. This is particularly useful in stability analysis, where one must bound the growth of perturbations in singular potentials. By establishing rigorous bounds on the energy, Hardy inequalities ensure that the physical or geometric models remain well-posed.

4.3. Computational Implications and Finite Element Methods

In numerical analysis, singularities pose severe challenges for standard discretization techniques. The finite element method (FEM) applied to singular elliptic problems suffers from reduced convergence rates if the mesh is not adapted to the singularity [7]. Weighted Hardy inequalities inform the design of adaptive mesh refinement (AMR) strategies. By utilizing a posteriori error estimators derived from Hardy-type bounds, computational algorithms can dynamically concentrate mesh elements near the singular points (e.g., the origin or the boundary).

Furthermore, these inequalities justify the use of graded meshes, ensuring that optimal convergence rates are recovered in the weighted Sobolev norms [5].

5. Illustrative Example

To concretize the classical Hardy inequality, consider the one-dimensional domain $(0,1)$ and the function $\mathbf{u}(x) = x(1 - x)$. Note that $\mathbf{u} \in H^{0,1}(0,1)$ since $\mathbf{u}(0) = \mathbf{u}(1) = 0$.

We compute the gradient (derivative) term: $\mathbf{u}'(x) = 1 - 2x$

$$\begin{aligned} \int_0^1 |\mathbf{u}'(x)|^2 dx &= \int_0^1 (1 - 2x)^2 dx = \int_0^1 (1 - 4x + 4x^2) dx = \left[x - 2x^2 + \frac{4}{3}x^3 \right]_0^1 \\ &= 1 - 2 + \frac{4}{3} = \frac{1}{3} \end{aligned}$$

Next, we compute the singular weighted term:

$$\int_0^1 \frac{|\mathbf{u}(x)|^2}{x^2} dx = \int_0^1 \frac{x^2(1 - x)^2}{x^2} dx = \int_0^1 (1 - x)^2 dx = \left[-\frac{(1 - x)^3}{3} \right]_0^1 = \frac{1}{3}$$

Applying the classical 1D Hardy inequality $\int_0^1 \frac{u^2}{x^2} dx \leq 4 \int_0^1 u'^2 dx$, we verify:

$$\frac{1}{3} \leq 4 \left(\frac{1}{3} \right) \Rightarrow \frac{1}{3} \leq \frac{4}{3}$$

The inequality holds strictly, illustrating how the gradient norm dominates the singular weighted norm, with the optimal constant 4 providing a strict upper bound [1, 2].

6. Discussion and Future Research

While the classical and weighted theories are well-established, several frontiers remain active areas of research:

Fractional Hardy Inequalities: With the rise of nonlocal models, extending Hardy inequalities to fractional Sobolev spaces $W^{s,p}$ is critical. Recent work focuses on fractional Hardy inequalities involving the fractional Laplacian $(-\Delta)^s$, which require novel techniques due to the nonlocal nature of the operator [3, 8].

Nonlinear p-Laplacian Problems: Generalizing from the linear Laplacian to the degenerate/singular p-Laplacian operator $\Delta_p \mathbf{u} = \nabla \cdot (|\nabla \mathbf{u}|^{p-2} \nabla \mathbf{u})$ introduces severe nonlinearities. Establishing sharp weighted Hardy inequalities for $p \neq 2$ remains a challenging endeavor, particularly in identifying the exact optimal constants for complex geometries [5].

Hardy Inequalities on Manifolds: Extending these inequalities to Riemannian manifolds with curvature constraints opens new avenues in geometric analysis. The interplay between the manifold's Ricci curvature and the Hardy constant is a rich area of study [3].

Data-Driven and Machine Learning Solvers: In computational mathematics, Physics-Informed Neural Networks (PINNs) and other machine learning-assisted PDE solvers struggle with singularities, often failing to converge or producing high errors near singular points. Future research will likely leverage weighted Hardy inequalities to design specialized loss functions or adaptive sampling strategies that penalize singular behavior, thereby stabilizing data-driven solvers.

7. Conclusion

Weighted Hardy inequalities are far more than abstract functional analytic curiosities; they are indispensable tools in the modern mathematical sciences. By providing rigorous, sharp estimates for functions with singular behavior, they ensure the well-posedness of singular elliptic problems and underpin the stability of variational formulations. Furthermore, their translation into the computational realm guides the development of robust, adaptive numerical schemes capable of resolving complex singularities. As mathematical models grow increasingly complex—incorporating fractional operators, nonlinearities, and data-driven paradigms—the theoretical and computational frameworks built upon weighted Hardy inequalities will remain vital to advancing both pure analysis and applied scientific computing.

References

- [1] Hardy, G. H., Littlewood, J. E., & Pólya, G. (1952). *Inequalities* (2nd ed.). Cambridge University Press.
- [2] Kufner, A., Maligranda, L., & Persson, L.-E. (2007). *The Hardy Inequality: About Its History and Some Related Results*. Vydavatelství VŠB.
- [3] Maz'ya, V. (2011). *Sobolev Spaces: With Applications to Elliptic Partial Differential Equations* (2nd ed.). Springer.
- [4] Adams, R. A., & Fournier, J. J. F. (2003). *Sobolev Spaces* (2nd ed.). Academic Press.
- [5] Edmunds, D. E., & Evans, W. D. (2004). *Hardy Operators, Function Spaces and Embeddings*. Springer.
- [6] Lieb, E. H., & Loss, M. (2001). *Analysis* (2nd ed.). American Mathematical Society (AMS).
- [7] Gilbarg, D., & Trudinger, N. S. (2001). *Elliptic Partial Differential Equations of Second Order*. Springer.
- [8] Ziemer, W. P. (1989). *Weakly Differentiable Functions: Sobolev Spaces and Functions of Bounded Variation*. Springer.

